

The following Listing of the Claims will replace all prior versions and all prior listings of the claims in the present application:

Listing of The Claims:

1. (Canceled)
2. (Canceled)
3. (Canceled)
4. (Previously Presented) The recombinant fusion polypeptide of claim 37, further comprising a third polypeptide peptide bonded to said recombinant fusion polypeptide.
5. (Original) The recombinant fusion polypeptide of claim 4 wherein said third polypeptide is a member of a specific binding pair.
6. (Original) The recombinant fusion polypeptide of claim 4 wherein said third polypeptide is fused to the amino terminus of said first polypeptide.
7. (Original) The recombinant fusion polypeptide of claim 4 wherein said third polypeptide is fused to the carboxy terminus of said second polypeptide sequence.
8. (Previously Presented) The recombinant fusion polypeptide of claim 37 wherein each of said first and said second polypeptide, independently, is a monomer of a multimeric protein selected from the group consisting of: *R. reniformis* GFP, *R. mulleri* GFP and *A. victoria* GFP.
9. (Previously Presented) The recombinant fusion polypeptide of claim 37 wherein both of said first and second polypeptides are monomers of a single multimeric protein selected from the group consisting of *R. reniformis* GFP, *R. mulleri* GFP and *A. victoria* GFP.
10. (Cancelled)
11. (Cancelled)
12. (Cancelled)

13. (Cancelled)

14. (Cancelled)

15. (Cancelled)

16. (Cancelled)

17. (Cancelled)

18. (Cancelled)

19. (Cancelled)

20. (Cancelled)

21. (Cancelled)

22. (Cancelled)

23. (Cancelled)

24. (Previously presented) A pair of polypeptides, one member of said pair comprising a polypeptide labeled with a fluorescent dye, and the other member of said pair comprising a recombinant fusion polypeptide, said fusion polypeptide comprising a first and a second polypeptide, wherein:

a) said first polypeptide is peptide bonded to said second polypeptide via a linker sequence,

b) said first and said second polypeptides are monomers of a multimeric fluorescent protein, and said first and said second polypeptides are not fluorescent donor and acceptor to each other; and

c) said recombinant fusion polypeptide is fluorescent when excited;

wherein said fluorescent dye and said recombinant fusion polypeptide are fluorescent donor and acceptor to each other.

25. (Currently Amended) A pair of recombinant fusion polypeptides comprising

a) a first fusion polypeptide comprising a first and a second polypeptide, wherein:

i) said first polypeptide is peptide bonded to said second polypeptide via a linker sequence,

ii) said first and said second polypeptides are monomers of a multimeric fluorescent protein, and said first and said second polypeptides are not fluorescent donor and acceptor to each other; and

iii) said recombinant fusion polypeptide is fluorescent when excited; and

b) a second fusion polypeptide comprising a first and a second polypeptide, wherein:

i) said first polypeptide is peptide bonded to said second polypeptide via a linker sequence,

ii) said first and said second polypeptides are monomers of a multimeric fluorescent protein, and said first and said second polypeptides are not fluorescent donor and acceptor to each other; and

iii) said recombinant fusion polypeptide is fluorescent when excited; wherein said first fusion polypeptide of (a) and said second fusion polypeptide of (b) are fluorescent donor and acceptor to each other.

26. (Original) The pair of recombinant fusion polypeptides of claim 25 wherein each of said first and second fusion polypeptides further comprises a third polypeptide, and wherein said third polypeptide of said first fusion polypeptide comprises a sequence which is different from said third polypeptide of said second fusion polypeptide.

27. (Cancelled)

28. (Cancelled)

29. (Cancelled)

30. (Original) A method of monitoring the interaction of two polypeptides of interest, said method comprising the steps of:

(a) contacting a first polypeptide and a second polypeptide wherein:

(i) said first polypeptide is a recombinant fusion polypeptide of claim 4 wherein said third polypeptide is a first polypeptide of interest;

(ii) said second polypeptide comprises a second polypeptide of interest and is fluorescently labeled; and

(iii) the fluorophores comprised by said first and second polypeptides are fluorescent donor and fluorescent acceptor to each other;

(b) exciting said donor fluorophore; and

(c) detecting fluorescent emission from said fluorescent acceptor, wherein said emission is indicative of the interaction of said first and said second polypeptides of interest.

31. (Original) The method of claim 30 wherein said second polypeptide comprises a fusion polypeptide of claim 5, wherein said third polypeptide of said second fusion polypeptide is different from said third polypeptide of said first fusion polypeptide.

32. (Original) The method of claim 30 wherein said contacting step is performed in vitro.

33. (Original) The method of claim 30 wherein said contacting step is performed in a cell.

34. (Original) The method of claim 33 wherein said contacting comprises the step of introducing nucleic acid encoding said first and said second polypeptides to a cell.

35. (Original) A method of screening for a compound that modulates the interaction of a first and a second member of a specific binding pair, said method comprising the steps of:

(a) contacting a first polypeptide and a second polypeptide in the presence and absence of a candidate modulator wherein:

(i) said first polypeptide is a recombinant fusion polypeptide of claim 5, wherein said member of a specific binding pair is said first member of a specific binding pair;

(ii) said second polypeptide is fluorescently labeled and comprises said second member of a specific binding pair; and

(iii) the fluorophores comprised by said first and second polypeptides are fluorescent donor and, acceptor to each other;

(b) exciting said donor fluorophore; and

(c) detecting the fluorescence of said acceptor fluorophore, wherein emission of the spectrum characteristic of said fluorescent acceptor indicates the interaction of said first and said second members of said specific binding pair, and wherein a change in said interaction in the presence of said candidate modulator indicates that said candidate modulator modulates the interaction of the members of said specific binding pair.

36. (Original) The method of claim 35 wherein said second polypeptide is a recombinant fusion polypeptide of claim 5 and said member of a specific binding pair comprised by said second polypeptide is said second member of a specific binding pair.

37. (Previously Presented) A recombinant fusion polypeptide comprising a first and a second polypeptide, wherein:

a) said first polypeptide is peptide bonded to said second polypeptide via a linker sequence of 5 to 50 amino acids in length,

b) said first and said second polypeptides are monomers of a multimeric fluorescent protein, and said first and said second polypeptides are not fluorescent donor and acceptor to each other; and

c) said recombinant fusion polypeptide is fluorescent when excited.

38. (Currently Amended) A recombinant fusion polypeptide comprising a first and a second polypeptide, wherein:

a) said first polypeptide is peptide bonded to said second polypeptide via a linker sequence selected from the group consisting of (Arg-Ala-Arg-Asp-Pro-Arg-Val-Pro-Val-Ala-Thr)₁₋₅ (SEQ ID NO: 8, 69-72), (Gly-Ser)₁₋₁₅ (SEQ ID NO: 9-23), (Thr-Ser-Pro)₁₋₁₅ (SEQ ID NO: 24-38), (Gly-Gly-Gly)₁₋₁₅ (SEQ ID NO: 39-53), (Glu-Lys)₁₋₁₅ (SEQ ID NO: 54-68), and (Gly₄Ser)₂₋₄ (SEQ ID NO: 5-7);

b) said first and said second polypeptides are monomers of a multimeric fluorescent protein, and said first and said second polypeptides are not fluorescent donor and acceptor to each other; and

c) said recombinant fusion polypeptide is fluorescent when excited.

39. (Cancelled)

40. (Cancelled)

41. (Cancelled)